



Life and Medical Sciences

Bilateral Ocular Hypertension (OHT) and Deep Vascularized Corneal Leukoma Following Measles: Why Should We get a Vaccine?

Kızamık Sonrası Bilateral Oküler Hipertansiyon (OHT) ve Derin Vaskülerize Korneal Lököm: Neden Aşı Yaptırmalıyız?

Mustafa KALAYCI¹ [ID]

¹Department of Ophthalmology, Mogadishu Somalia-Turkey Recep Tayyip Erdoğan Training and Research Hospital, Mogadishu, Somalia.

Article Info: Received; 23.12.2019. Accepted; 03.01.2020. Published; 03.01.2020.

Correspondence: Mustafa Kalaycı; MD, Department of Ophthalmology, Mogadishu Somalia-Turkey Recep Tayyip Erdoğan Training and Research Hospital, Mogadishu, Somalia. E-mail: drkalayci07@hotmail.com

Cite as: Kalaycı M. Bilateral Ocular Hypertension (OHT) and Deep Vascularized Corneal Leukoma Following Measles: Why Should We get a Vaccine? Life Med Sci 2022; 1(1): 24-26.

Abstract

Measles is characterized by a pronounced exanthema, pathognomonic enanthema accompanied by cough, fever, and conjunctivitis. A 2-year-old girl was presented to ophthalmology department by her family with whitening and vision loss in her eyes in Mogadishu - Somalia. Retinal and vitreous layers were normal in both eyes on B-scan ultrasound. Intraocular pressure was 34 mmHg bilaterally. Bilateral ocular hypertension and deep vascularized corneal leukoma were diagnosed. This is the first case in the literature that developed bilateral ocular hypertension and deep vascularized leukoma after measles. Children are more susceptible to the severity and side effects of measles, especially in infancy. We recommend vaccination to prevent post-disease complications in young children.

Keywords: Somalia, Keratoplasty, Conjunctivitis, Ocular ultrasound.

Özet

Kızamık, belirgin bir ekzantem ve patognomonik enantem olarak adlandırılan öksürük, ateş, konjunktivit ile karakterize bir hastalıktır. 2 yaşında bir kız çocuğu Mogadişu - Somali'de gözlerinde beyaz renkte değişim ve görme kaybı ile birlikte oftalmoloji bölümüne ailesi tarafından getirildi. B-tarama ultrasonunda her iki gözde retina ve vitreus yapıları normaldi. Göz içi basıncı her iki gözde 34 idi. Bilateral lököm ve oküler hipertansiyon tanısı konuldu. Yapılan literatür taramasında kızamık hastalığı sonrası bilateral lököm ve oküler hipertansiyon birlikteliği olan ilk vaka olduğu saptandı. Çocuklar, özellikle bebeklik döneminde kızamıkların ciddiyetine ve yan etkilerine daha duyarlıdır. Küçük çocuklarda hastalık sonrası komplikasyonları önlemek için aşı yapılmasını öneririz.

Anahtar Kelimeler: Somali, Keratoplasti, Konjunktivit, Oküler ultrason.

Introduction

Measles is a highly contagious acute viral disease characterized by a pronounced exanthema, pathognomonic enanthema accompanied by cough, fever and conjunctivitis [1]. Measles was a leading global cause of morbidity and mortality in children prior to vaccination. It was responsible for more than

2 million deaths a year. In the 1980s, global measles vaccination under the leadership of the Expanded Immunization Program has rapidly lowered this rate worldwide [2]. However, the vaccination program is still not fully implemented in less developed countries.

Case Report

A 2-year-old girl was presented to ophthalmology department by her family with whitening and vision loss in her eyes in Mogadishu - Somalia. The patient had never been examined by an ophthalmologist before. Her family explained that the patient had measles 6 months ago and that she rubbed his eyes with his finger after the rash. The family said the patient had never been vaccinated against measles (Figure 1).



Figure 1: Bilateral ocular hypertension and corneal leukoma of the patient.




Figure 2: Ocular USG of the patient.

Retinal and vitreous layers were normal in both eyes on B-scan ultrasound (Figure 2). In the examination performed under general anesthesia, intraocular pressure was measured by Schiottz indentation tonometer and it was 34 mmHg bilaterally. After ultrasonic pachymetry measurement, the corneal central thickness of the right eye was 620 μm and the left eye was 634 μm . Glaucoma could not be diagnosed because the iridocorneal angle and optic disc could not be evaluated. The patient had superficial and deep vascularization in both corneas. Bilateral ocular hypertension and corneal leukoma were diagnosed. No other systemic or ophthalmic disease was detected in the examination and laboratory results of the patient. 2-year-old patient was asked to be referred for keratoplasty procedures because of rapid amblyopia. However, the patient's family did not accept surgery Brinzolamide + timolol and travoprost eye drops were started to reduce OHT. At 1 week follow-up, eye pressure was 14 mmHg. According to our research this is the first case in the literature that developed bilateral OHT and deep vascularized leukoma after measles.

Discussion

Children are more susceptible to the severity and side effects of measles, especially in infancy [3,4]. In measles disease, ocular findings are associated with acute keratoconjunctivitis and usually heal without sequelae [5]. Although few cases of bilateral corneal leukoma after measles have been previously reported, the fact that bilateral ocular hypertension and corneal leukoma coexist, suggests that measles may play a role in the etiopathogenesis of OHT and progression to glaucoma. Antibodies formed in the body after measles may cause iridocyclitis in both anterior chambers and cause angle closure.

Infancy, very severe lower respiratory tract, malnutrition, and immunosuppression are the major risk factors that increase morbidity and mortality for measles [6]. In underdeveloped countries such as Somalia, where malnutrition is common, measles complications can be severe in terms of the eye as a result of the lack of measles vaccination. The consequences of these problems can lead to blindness [7].

Conclusion

Despite the presence of a safe and effective vaccine, measles leading death among young children around the world is one of the reasons. Measles of vaccination campaigns mortality has been observed to have a significant effect. Today, the tendency of families not to have their children measles vaccine

is increasing all over the world. We recommend vaccination to prevent post-disease complications in young children.

Declaration of interest: The author declares no conflict of interest and alone is responsible for the content and writing of the paper.

This article previously published as: "Somalia Turkey Journal of Medical Science 2020; 1(1): 9-10." Currently, Somalia Turkey Journal of Medical Science was merged with Life and Medical Sciences.

References

1. Rota PA, Moss WJ, Takeda M, de Swart RL, Thompson KM, Goodson JL. Measles. *Nat Rev Dis Primers* 2016; 2: 16049. [[Crossref](#)] [[PubMed](#)]
2. Moss WJ. Measles. *Lancet* 2017; 390(10111): 2490-502. [[Crossref](#)] [[PubMed](#)]
3. Cherry JD, Zahn M. Clinical Characteristics of Measles in Previously Vaccinated and Unvaccinated Patients in California. *Clin Infect Dis* 2018; 67(9): 1315-9. [[Crossref](#)] [[PubMed](#)]
4. Sauter JJ. Why measles makes so many children blind. *Trop Doct* 1982; 12(4 Pt 2): 219-22. [[Crossref](#)] [[PubMed](#)]
5. Yoshikura H. Impact of population size on incidence of rubella and measles in comparison with that of other infectious diseases [published correction appears in *Jpn J Infect Dis.* 2015; 68(1): 80]. *Jpn J Infect Dis.* 2014; 67(6): 447-57. [[Crossref](#)] [[PubMed](#)]
6. Hirfanoğlu T, Tanir G, Karacan C, Göl N. Clinical characteristics, complications and prognosis of seventy-nine measles cases. *Journal of Ankara University Faculty of Medicine* 2006; 59(3): 98-103.
7. Mado SM, Idris HW, Abdullahi SM, Musa S. Anterior staphyloma in a 6-month-old, severely malnourished child following measles: A case report. *Sub-Saharan African Journal of Medicine* 2018; 5(1): 29-31. [[Crossref](#)]